

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: GREENE, ET AL.

(ii) TITLE OF INVENTION: Human Tumor Necrosis Factor
Receptor

(iii) NUMBER OF SEQUENCES: 9

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: CARELLA, BYRNE, BAIN, GILFILLAN,
CECCHI, STEWART & OLSTEIN

(B) STREET: 6 BECKER FARM ROAD

(C) CITY: ROSELAND

(D) STATE: NEW JERSEY

(E) COUNTRY: USA

(F) ZIP: 07068

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 3.5 INCH DISKETTE

(B) COMPUTER: IBM PS/2

(C) OPERATING SYSTEM: MS-DOS

(D) SOFTWARE: WORD PERFECT 5.1

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE: Concurrently

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA

(A) APPLICATION NUMBER: PCT/US95/03216

(B) FILING DATE: 15 MAR 1995

*sub
a'*

(viii) ATTORNEY/AGENT INFORMATION:
 (A) NAME: FERRARO, GREGORY D.
 (B) REGISTRATION NUMBER: 36,134
 (C) REFERENCE/DOCKET NUMBER: 325800-381

(ix) TELECOMMUNICATION INFORMATION:
 (A) TELEPHONE: 201-994-1700
 (B) TELEFAX: 201-994-1744

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 1173 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAACAAGT	TGCTGTGCTG	CGCGCTCGTG	TTTCTGGACA	TCTCCATTAA	GTGGACCACC	60
CAGGAAACGT	TTCCTCCAAA	GTACCTTCAT	TATGACGAAG	AAACCTCTCA	TCAGCTGTTG	120
TGTGACAAAT	GTCCTCCTGG	TACCTACCTA	AAACAACACT	GTACAGCAAA	GTGGAAGACC	180
GTGTGCGCCC	CTTGCCCTGA	CCACTACTAC	ACAGACAGCT	GGCACACCAG	TGACGAGTGT	240
CTATACTGCA	GCCCCGTGTG	CAAGGAGCTG	CAGTACGTCA	AGCAGGAGTG	CAATCGCACC	300
CACAACCGCG	TGTGCGAATG	CAAGGAAGGG	CGCTACCTTG	AGATAGAGTT	CTGCTTGAAA	360
CATAGGAGCT	GCCCTCCTGG	ATTTGGAGTG	GTGCAAGCTG	GAACCCGAGA	GCGAAATACA	420
GTTTGCAAAA	GATGTCCAGA	TGGGTTCTTC	TCAAATGAGA	CGTCATCTAA	AGCACCCTGT	480
AGAAAACACA	CAAATTGCAG	TGTCTTTGGT	CTCCTGCTAA	CTCAGAAAGG	AAATGCAACA	540
CACGACAACA	TATGTTCCGG	AAACAGTGAA	TCAACTCAAA	AATGTGGAAT	AGATGTTACC	600
CTGTGTGAGG	AGGCATTCTT	CAGGTTTGCT	GTTCTTACAA	AGTTTACGCC	TAACTGGCTT	660
AGTGTCTTTG	TAGACAATTT	GCCTGGCACC	AAAGTAAACG	CAGAGAGTGT	AGAGAGGATA	720
AAACGGCAAC	ACAGCTCACA	AGAACAGACT	TTCCAGCTGC	TGAAGTTATG	GAAACATCAA	780
AACAAAGACC	AAGATATAGT	CAAGAAGATC	ATCCAAGATA	TTGACCTCTG	TGAAAACAGC	840
GTGCAGCGGC	ACATTGGACA	TGCTAACCTC	ACCTTCGAGC	AGCTTCGTAG	CTTGATGGAA	900
AGCTTACCGG	GAAAGAAAGT	GGGAGCAGAA	GACATTGAAA	AAACAATAAA	GGCATGCAAA	960
CCCAGTGACC	AGATCCTGAA	GCTGCTCAGT	TTGTGGCGAA	TAAAAAATGG	CGACCAAGAC	1020

ACCTTGAAGG GCCTAATGCA CGCACTAAAG CACTCAAAGA CGTACCACTT TCCCACAAAC 1080
 TGTCACTCAG AGTCTAAAGA AGACCATCAG GTTCCTTCAC AGCTTCACAA TGTACAAATT 1140
 GTATCAGAAG TTATTTTGTAG AAATGATAGG TAA 1173

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 390 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Asn	Lys	Leu	Leu	Cys	Cys	Ala	Leu	Val	Phe	Leu	Asp	Ile	Ser	
	-20					-15					-10				
Ile	Lys	Trp	Thr	Thr	Gln	Glu	Thr	Phe	Pro	Pro	Lys	Tyr	Leu	His	
	-5					1					5				
Tyr	Asp	Glu	Glu	Thr	Ser	His	Gln	Leu	Leu	Cys	Asp	Lys	Cys	Pro	
	10					15					20				
Pro	Gly	Thr	Tyr	Leu	Lys	Gln	His	Cys	Thr	Ala	Lys	Trp	Lys	Thr	
	25					30					35				
Val	Cys	Ala	Pro	Cys	Pro	Asp	His	Tyr	Tyr	Thr	Asp	Ser	Trp	His	
	40					45					50				
Thr	Ser	Asp	Glu	Cys	Leu	Tyr	Cys	Ser	Pro	Val	Cys	Lys	Glu	Leu	
	55					60					65				
Gln	Tyr	Val	Lys	Gln	Glu	Cys	Asn	Arg	Thr	His	Asn	Arg	Val	Cys	
	70					75					80				
Glu	Cys	Lys	Glu	Gly	Arg	Tyr	Leu	Glu	Ile	Glu	Phe	Cys	Leu	Lys	
	85					90					95				
His	Arg	Ser	Cys	Pro	Pro	Gly	Phe	Gly	Val	Val	Gln	Ala	Gly	Thr	
	100					105					110				
Pro	Glu	Arg	Asn	Thr	Val	Cys	Lys	Arg	Cys	Pro	Asp	Gly	Phe	Phe	
	115					120					125				

Ser	Asn	Glu	Thr	Ser	Ser	Lys	Ala	Pro	Cys	Arg	Lys	His	Thr	Asn
130						135				140				
Cys	Ser	Val	Phe	Gly	Leu	Leu	Leu	Thr	Gln	Lys	Gly	Asn	Ala	Thr
145						150				155				
His	Asp	Asn	Ile	Cys	Ser	Gly	Asn	Ser	Glu	Ser	Thr	Gln	Lys	Cys
160						165				170				
Gly	Ile	Asp	Val	Thr	Leu	Cys	Glu	Glu	Ala	Phe	Phe	Arg	Phe	Ala
175						180				185				
Val	Pro	Thr	Lys	Phe	Thr	Pro	Asn	Trp	Leu	Ser	Val	Leu	Val	Asp
190						195				200				
Asn	Leu	Pro	Gly	Thr	Lys	Val	Asn	Ala	Glu	Ser	Val	Glu	Arg	Ile
205						210				215				
Lys	Arg	Gln	His	Ser	Ser	Gln	Glu	Gln	Thr	Phe	Gln	Leu	Leu	Lys
220						225				230				
Leu	Trp	Lys	His	Gln	Asn	Lys	Asp	Gln	Asp	Ile	Val	Lys	Lys	Ile
235						240				245				
Ile	Gln	Asp	Ile	Asp	Leu	Cys	Glu	Asn	Ser	Val	Gln	Arg	His	Ile
250						255				260				
Gly	His	Ala	Asn	Leu	Thr	Phe	Glu	Gln	Leu	Arg	Ser	Leu	Met	Glu
265						270				275				
Ser	Leu	Pro	Gly	Lys	Lys	Val	Gly	Ala	Glu	Asp	Ile	Glu	Lys	Thr
280						285				290				
Ile	Lys	Ala	Cys	Lys	Pro	Ser	Asp	Gln	Ile	Leu	Lys	Leu	Leu	Ser
295						300				305				
Leu	Trp	Arg	Ile	Lys	Asn	Gly	Asp	Gln	Asp	Thr	Leu	Lys	Gly	Leu
310						315				320				
Met	His	Ala	Leu	Lys	His	Ser	Lys	Thr	Tyr	His	Phe	Pro	Thr	Asn
325						330				335				
Cys	His	Ser	Glu	Ser	Lys	Glu	Asp	His	Gln	Val	Pro	Ser	Gln	Leu
340						345				350				
His	Asn	Val	Gln	Ile	Val	Ser	Glu	Val	Ile	Phe	Arg	Asn	Asp	Arg
355						360				365				

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 33 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCCAGAGGAT CCGAAACGTT TCCTCCAAAG TAC

33

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 33 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGGCTTCTAG AATTACCTAT CATTTCTAAA AAT

33

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 31 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCGCGGATCC ATGAACAAGT TGCTGTGCTG C

31

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 34 BASE PAIRS

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCGCTCTAGA TTACCTATCA TTTCTAAAAA TAAC

34

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 31 BASE PAIRS

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GCGCGGTACC TCAGTGGTTT GGGCTCCTCC C

31

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 39 BASE PAIRS
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCCAGAGGAT CCGCCACCAT GAACAAGTTG CTGTGCTGC

39

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 60 BASE PAIRS
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGGCTTCTAG AATCAAGCGT AGTCTGGGAC GTCGTATGGG TACCTATCAT TTCTAAAAAT

60

6. A process for producing a polypeptide comprising: expressing from the host cell of Claim 5 the polypeptide encoded by said DNA.

7. A process for producing cells capable of expressing a polypeptide comprising transforming or transfecting the cells with the vector of Claim 4.

8. A receptor polypeptide comprising a member selected from the group consisting of:

(i) a polypeptide having the deduced amino acid sequence of SEQ ID NO:2 and fragments, analogs and derivatives thereof; and

(ii) a polypeptide encoded by the cDNA of ATCC Deposit No. 75899 and fragments, analogs and derivatives of said polypeptide.

9. An antibody against the polypeptide of claim 8.

10. A compound which activates the polypeptide of claim 8.

11. A compound which inhibits activation the polypeptide of claim 8.

12. A method for the treatment of a patient having need to activate a TNF receptor comprising: administering to the patient a therapeutically effective amount of the compound of claim 10.

13. A method for the treatment of a patient having need to inhibit a TNF receptor comprising: administering to the patient a therapeutically effective amount of the compound of claim 11.

14. The method of claim 12 wherein said compound is a polypeptide and a therapeutically effective amount of the compound is administered by providing to the patient DNA encoding said agonist and expressing said agonist *in vivo*.

15. The method of claim 13 wherein said compound is a polypeptide and a therapeutically effective amount of the compound is administered by providing to the patient DNA encoding said antagonist and expressing said antagonist *in vivo*.

16. A method for identifying compounds which bind to and activate the receptor polypeptide of claim 8 comprising:

contacting a cell expressing on the surface thereof the receptor polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor polypeptide, with a compound under conditions sufficient to permit binding of the compound to the receptor polypeptide; and

identifying if the compound is capable of receptor binding by detecting the signal produced by said second component.

17. A method for identifying compounds which bind to and inhibit activation of the polypeptide of claim 8 comprising:

contacting a cell expressing on the surface thereof the receptor polypeptide, said receptor being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said receptor polypeptide, with an analytically detectable ligand known to bind to the receptor polypeptide

and a compound to be screened under conditions to permit binding to the receptor polypeptide; and

determining whether the compound inhibits activation of the polypeptide by detecting the absence of a signal generated from the interaction of the ligand with the polypeptide.

18. A process for diagnosing a disease or a susceptibility to a disease related to an under-expression of the polypeptide of claim 8 comprising:

determining a mutation in the nucleic acid sequence encoding said polypeptide.

19. The polypeptide of Claim 8 wherein the polypeptide is a soluble fragment of the polypeptide and is capable of binding a ligand for the receptor.

20. A diagnostic process comprising:
analyzing for the presence of the polypeptide of claim 19 in a sample derived from a host.

Added B8